

The Cholangiocarcinoma Treatment Landscape: Expanding Horizons and Improving Outcomes (Recorded Podcast)

REFERENCES

1. National Comprehensive Cancer Network (NCCN). Hepatobiliary cancers. Version 5.2020. https://www.nccn.org/professionals/physician_gls/pdf/hepatobiliary.pdf. Accessed November 20,2020.
2. Banales JM, Marin JJG, Lamarca A, et al. Cholangiocarcinoma 2020: the next horizon in mechanisms and management. *Nat Rev Gastroenterol Hepatol*. 2020;17(9):557-588.
3. Primrose JN, Fox RP, Palmer DH, et al. Capecitabine compared with observation in resected biliary tract cancer (BILCAP): a randomised, controlled, multicentre, phase 3 study. *Lancet Oncol*. 2019;20:663-673.
4. Valle J, Wasan H, Palmer DH, et al. Cisplatin plus gemcitabine versus gemcitabine for biliary tract cancer. *N Eng J Med*. 2010;362(14):1273-1281.
5. Shroff RT, Javle MM, Xiao L, et al. Gemcitabine, cisplatin, and nab-paclitaxel for the treatment of advanced biliary tract cancers: a phase 2 clinical trial. *JAMA Oncol*. 2019;5(6):824-830.
6. Tella SH, Kommalapati A, Borad MJ, Mahipal A. Second-line therapies in advanced biliary tract cancers. *Lancet Oncol*. 2020;21(1):e29-e41.
7. Abou-Alfa GK, Macarulla T, Javle MM, et al. Ivosidenib in IDH1-mutant, chemotherapy-refractory cholangiocarcinoma (ClarIDHy): a multicentre, randomised, double-blind, placebo-controlled, phase 3 study. *Lancet Oncol*. 2020;21(6):796-807.
8. About-Alfa GK, Sahai V, Hollebecque A, et al. Pemigatinib for previously treated, locally advanced or metastatic cholangiocarcinoma: a multicentre, open-label, phase 2 study. *Lancet Oncol*. 2020;21(5):671-684.
9. Mahipal A, Tella SH, Kommalapati A, et al. Prevention and treatment of FGFR inhibitor-associated toxicities. *Crit Rev Oncol Hematol*. 2020;155:103091.